



Acetaminophen

Drops, Suspension, various tablet forms, suppositories, in combination with other drugs

Usages:

- Fever reliever
- Pain reliever (common colds, flu, headaches, sore throat, immunizations, toothaches, ear pain , and others)

Side Effects:

- None unless in an overdose situation. Do not exceed the recommended dose
- Rare sensitivity reaction do occur and the drug should not be given

Warnings/ Do not use:

- With any other product containing acetaminophen
- For more than 3 days for fever unless directed by a doctor
- For more than 5 days for pain unless directed by a doctor
- Symptoms do not improve
- New symptoms occur
- Pain of fever persists or gets worse
Redness of swelling is present
- Sore throat is severe, lasts for more than 2 days or occurs with fever, headache, rash, nausea, vomiting . Consult doctor.
- Do not use if plastic carton wrap/ or bottle wrap imprinted “Safety Seal@” is broken or missing.
- Do not use if carton is opened or if neck wrap or foil inner seal imprinted “Safety Seal@” is broken or missing.
- Do not use if carton is opened or if blister unit is broken
- Phenylketonuric people should not use as various flavors contain different amounts of phenylketouria

Abuse/Overdose:

- Get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.
- Overdose may occur following ingestion of greater 7.5 grams over a period of 8 hours and adults and adolescents.
- Overdose in children < 12 years follows ingestion of 150mg/kg